

# Off-Diagonal Long-Range Order: Meissner Effect and Flux Quantization

H.T. Nieh and Gang Su\*

Institute for Theoretical Physics

State University of New York at Stony Brook

Stony Brook, New York 11794, USA

Bao-Heng Zhao

Department of Physics, Graduate School

Chinese Academy of Sciences, P. O. Box 3908

Beijing 100039, China

## Abstract

There has been a proof by Sewell that the hypothesis of off-diagonal long-range order in the reduced density matrix  $\rho_2$  implies the Meissner effect. We present in this note an elementary and straightforward proof that not only the Meissner effect but also the property of magnetic flux quantization follows from the hypothesis. It is explicitly shown that the two phenomena are closely related, and phase coherence is the origin for both.

PACS numbers: 05.30.-d; 74.20.-z; 74.25.Ha

---

\*On leave of absence from the Graduate School, Chinese Academy of Sciences, Beijing, China.

## Introduction

Penrose and Onsager<sup>1</sup> first proposed that the mathematical statement of Bose-Einstein condensation in the case of interacting boson systems is the existence of off-diagonal long-range order (ODLRO) in the reduced density matrix  $\rho_1$ . Yang<sup>2</sup> later made a thorough study of the subject and generalized the concept to interacting fermion systems. It is now generally accepted that the superconducting state of electrons is characterized by the existence of off-diagonal long-range order in the reduced density matrix  $\rho_2$ .

In view of the fundamental importance of the concept of ODLRO, it is important that there be a rigorous proof of the assertion that ODLRO in the reduced density matrix  $\rho_2$  implies both the Meissner effect and the quantization of magnetic flux, which are the basic characteristic properties of superconducting states. There has been a recent proof by Sewell<sup>3</sup> of the Meissner effect as a consequence of ODLRO, under certain simplifying assumptions. We shall extend in this note the proof to the case of magnetic flux quantization.

The proof we present is simple and elementary, and clearly shows that both the Meissner effect and the flux quantization are closely related phenomena, and the underlying origin for both is the coherence of phases.

## System in Uniform Magnetic Field

We consider a system of  $N$  electrons inside a crystal, which is placed in a uniform external magnetic field. We make the simplifying assumption that the magnetic field  $\mathbf{B}$  inside the material is also uniform, with the corresponding vector potential, given by  $\mathbf{A}(\mathbf{r})$ , which is of the general form:

$$\mathbf{A}(\mathbf{r}) = \mathbf{A}_0(\mathbf{r}) + \nabla\phi(\mathbf{r}), \quad (1)$$

where  $\mathbf{A}_0(\mathbf{r}) = \frac{1}{2}\mathbf{B} \times \mathbf{r}$ , and the gauge  $\phi(\mathbf{r})$  is, in general, multi-valued in  $\mathbf{r}$ . The Hamiltonian for the system is

$$H = \sum_j \frac{1}{2m} \left[ \frac{\hbar}{i} \nabla_j + \frac{e}{c} \mathbf{A}(\mathbf{r}_j) \right]^2 + \frac{1}{2} \sum_{j,k} V(r_{jk}) + \sum_j V_c(\mathbf{r}_j), \quad (2)$$

where  $V_c(\mathbf{r}_j)$  represents the interaction of the crystal lattice on the  $j$ -th electron, and the two-particle potential  $V(r_{jk})$ , which includes the effective interaction due to phonon exchange,

depends on the distance between the two electrons. Let

$$\psi_n(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N), \quad n = 1, 2, \dots,$$

represent a complete orthonormal set of single-valued and properly antisymmetrized eigenfunctions of the Hamiltonian

$$H\psi_n(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N) = E_n\psi_n(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N). \quad (3)$$

We neglect the edge effects near the boundaries, and assume the usual periodic boundary conditions for the wave functions. The phenomenon of flux penetration near boundaries is beyond the scope of the present idealized considerations.

## Two Complete Sets of Eigenfunctions

Now, consider a space displacement

$$\mathbf{r}_j \rightarrow \mathbf{r}_j - \mathbf{a},$$

under which

$$V(r_{jk}) \rightarrow V(r_{jk}),$$

$$\mathbf{A}(\mathbf{r}_j) \rightarrow \mathbf{A}(\mathbf{r}_j - \mathbf{a}) = \mathbf{A}(\mathbf{r}_j) + \nabla_j[\mathbf{a} \cdot \mathbf{A}_0(\mathbf{r}_j) + \phi(\mathbf{r}_j - \mathbf{a}) - \phi(\mathbf{r}_j)]. \quad (4)$$

We observe that a space displacement induces a gauge transformation in the vector potential. It is this property that makes it possible for a simple proof of the flux quantization and the Meissner effect as consequences of the ODLRO hypothesis. The lattice potential  $V_c(\mathbf{r}_j)$  has the symmetry of the lattice, and is invariant under a displacement only if the displacement respects the symmetry of the crystal lattice.

We first consider the case where the lattice potential  $V_c(\mathbf{r}_j)$  in the Hamiltonian is omitted, and will come back later to consider the case where it is not. Without the lattice potential, the eigen-equation (3), under the displacement  $\mathbf{r}_j \rightarrow \mathbf{r}_j - \mathbf{a}$ , becomes

$$\begin{aligned} & \left\{ \sum_j \frac{1}{2m} \left[ \frac{\hbar}{i} \nabla_j + \frac{e}{c} \mathbf{A}(\mathbf{r}_j - \mathbf{a}) \right]^2 + \frac{1}{2} \sum_{j,k} V(r_{jk}) \right\} \psi_n(\mathbf{r}_1 - \mathbf{a}, \mathbf{r}_2 - \mathbf{a}, \dots, \mathbf{r}_N - \mathbf{a}) \\ & = E_n \psi_n(\mathbf{r}_1 - \mathbf{a}, \mathbf{r}_2 - \mathbf{a}, \dots, \mathbf{r}_N - \mathbf{a}). \end{aligned} \quad (5)$$

Noting that a displacement induces a gauge transformation, as in (4), in the vector potential, we can absorb the gauge into a phase for the electron wave function to obtain

$$\begin{aligned} & \left\{ \sum_j \frac{1}{2m} \left[ \frac{\hbar}{i} \nabla_j + \frac{e}{c} \mathbf{A}(\mathbf{r}_j) \right]^2 + \frac{1}{2} \sum_{j,k} V(r_{jk}) \right\} \exp \left[ \frac{ie}{c\hbar} \sum_j \chi_{\mathbf{a}}(\mathbf{r}_j) \right] \psi_n(\mathbf{r}_1 - \mathbf{a}, \mathbf{r}_2 - \mathbf{a}, \dots, \mathbf{r}_N - \mathbf{a}) \\ &= E_n \exp \left[ \frac{ie}{c\hbar} \sum_j \chi_{\mathbf{a}}(\mathbf{r}_j) \right] \psi_n(\mathbf{r}_1 - \mathbf{a}, \mathbf{r}_2 - \mathbf{a}, \dots, \mathbf{r}_N - \mathbf{a}), \end{aligned} \quad (6)$$

where

$$\chi_{\mathbf{a}}(\mathbf{r}_j) = \mathbf{a} \cdot \mathbf{A}_0(\mathbf{r}_j) + \phi(\mathbf{r}_j - \mathbf{a}) - \phi(\mathbf{r}_j). \quad (7)$$

It is seen that  $\psi'_n(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N)$ , given by

$$\psi'_n(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N) = \exp \left[ \frac{ie}{c\hbar} \sum_j \chi_{\mathbf{a}}(\mathbf{r}_j) \right] \psi_n(\mathbf{r}_1 - \mathbf{a}, \mathbf{r}_2 - \mathbf{a}, \dots, \mathbf{r}_N - \mathbf{a}), \quad (8)$$

are again eigenfunctions of the original Hamiltonian. In the following we will only consider the case of  $\mathbf{a}$  to be infinitesimal on macroscopic scale<sup>4</sup>. In this case the phase  $\chi_{\mathbf{a}}(\mathbf{r})$  is single-valued in  $\mathbf{r}$ . Therefore,  $\psi'_n$  form another complete orthonormal set of single-valued eigenfunctions. We thus have two bases of eigenfunctions,  $\psi_n$  and  $\psi'_n$ .

### Reduced Density Matrix for Two Particles

The reduced density matrix element  $\rho_2(\mathbf{r}'_1, \mathbf{r}'_2; \mathbf{r}_1, \mathbf{r}_2)$  is defined in terms of a complete orthonormal set of energy eigenfunctions by

$$\begin{aligned} \rho_2(\mathbf{r}'_1, \mathbf{r}'_2; \mathbf{r}_1, \mathbf{r}_2) &= \int \cdots \int \frac{d\mathbf{r}_3 \cdots d\mathbf{r}_N}{(N-2)!} \frac{1}{Z} \sum_n \exp(-E_n/kT) \\ &\quad \times \psi_n(\mathbf{r}'_1, \mathbf{r}'_2, \mathbf{r}_3, \dots, \mathbf{r}_N) \psi_n^*(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \dots, \mathbf{r}_N), \end{aligned} \quad (9)$$

where  $Z$  is the partition function, and the normalization is such that

$$\text{Tr} \rho_2 = \int \int \rho_2(\mathbf{r}_1, \mathbf{r}_2; \mathbf{r}_1, \mathbf{r}_2) d\mathbf{r}_1 d\mathbf{r}_2 = N(N-1).$$

We note that  $\rho_2(\mathbf{r}'_1, \mathbf{r}'_2; \mathbf{r}_1, \mathbf{r}_2)$  is single-valued, as the wave-functions are single-valued. Since the reduced density matrices are basis independent, we can equally represent  $\rho_2(\mathbf{r}'_1, \mathbf{r}'_2; \mathbf{r}_1, \mathbf{r}_2)$  in terms of the primed set of energy eigenfunctions:

$$\rho_2(\mathbf{r}'_1, \mathbf{r}'_2; \mathbf{r}_1, \mathbf{r}_2) = \int \cdots \int \frac{d\mathbf{r}_3 \cdots d\mathbf{r}_N}{(N-2)!} \frac{1}{Z} \sum_n \exp(-E_n/kT)$$

$$\times \psi'_n(\mathbf{r}'_1, \mathbf{r}'_2, \mathbf{r}_3, \dots, \mathbf{r}_N) \psi_n'^*(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \dots, \mathbf{r}_N), \quad (10)$$

Expressing the primed eigenfunctions in the above equation in terms of the original eigenfunctions, we obtain

$$\begin{aligned} \rho_2(\mathbf{r}'_1, \mathbf{r}'_2; \mathbf{r}_1, \mathbf{r}_2) &= \exp\left\{\frac{ie}{c\hbar}[\chi_{\mathbf{a}}(\mathbf{r}'_1) + \chi_{\mathbf{a}}(\mathbf{r}'_2) - \chi_{\mathbf{a}}(\mathbf{r}_1) - \chi_{\mathbf{a}}(\mathbf{r}_2)]\right\} \\ &\times \int \dots \int \frac{d\mathbf{r}_3 \dots d\mathbf{r}_N}{(N-2)!} \frac{1}{Z} \sum_n \exp(-E_n/kT) \\ &\times \psi_n(\mathbf{r}'_1 - \mathbf{a}, \mathbf{r}'_2 - \mathbf{a}, \mathbf{r}_3 - \mathbf{a}, \dots, \mathbf{r}_N - \mathbf{a}) \psi_n^*(\mathbf{r}_1 - \mathbf{a}, \mathbf{r}_2 - \mathbf{a}, \mathbf{r}_3 - \mathbf{a}, \dots, \mathbf{r}_N - \mathbf{a}). \end{aligned}$$

On account of the periodic boundary conditions satisfied by the wave functions, we can make an innocent shift of the integration variables in the integral in the above equation, resulting in the relation of the density matrix at different space points:

$$\rho_2(\mathbf{r}'_1, \mathbf{r}'_2; \mathbf{r}_1, \mathbf{r}_2) = \exp\left\{\frac{ie}{c\hbar}[\chi_{\mathbf{a}}(\mathbf{r}'_1) + \chi_{\mathbf{a}}(\mathbf{r}'_2) - \chi_{\mathbf{a}}(\mathbf{r}_1) - \chi_{\mathbf{a}}(\mathbf{r}_2)]\right\} \rho_2(\mathbf{r}'_1 - \mathbf{a}, \mathbf{r}'_2 - \mathbf{a}; \mathbf{r}_1 - \mathbf{a}, \mathbf{r}_2 - \mathbf{a}). \quad (11)$$

This equation, as we shall see, is of basic importance.

### Off-Diagonal Long-Range Order

We can represent the reduced density matrix element in the spectral form:

$$\rho_2(\mathbf{r}'_1, \mathbf{r}'_2; \mathbf{r}_1, \mathbf{r}_2) = \sum_s \alpha_s \Phi_s(\mathbf{r}'_1, \mathbf{r}'_2) \Phi_s^*(\mathbf{r}_1, \mathbf{r}_2), \quad (12)$$

where  $\alpha_s$  are eigenvalues, and  $\Phi_s(\mathbf{r}_1, \mathbf{r}_2)$  the properly normalized eigenfunctions of  $\rho_2$ :

$$\int \int \Phi_s^*(\mathbf{r}_1, \mathbf{r}_2) \Phi_s(\mathbf{r}_1, \mathbf{r}_2) d\mathbf{r}_1 d\mathbf{r}_2 = 1.$$

We are interested in the property of  $\rho_2$  in the off-diagonal long-range (ODLR) limit:

$$|\mathbf{r}'_j - \mathbf{r}_k| \rightarrow \infty, \quad j, k = 1, 2$$

while keeping  $|\mathbf{r}_1 - \mathbf{r}_2|$  and  $|\mathbf{r}'_1 - \mathbf{r}'_2|$  finite. The hypothesis of ODLRO in  $\rho_2$  is the statement that in the ODLR limit the reduced density matrix element factorizes in the form

$$\rho_2(\mathbf{r}'_1, \mathbf{r}'_2; \mathbf{r}_1, \mathbf{r}_2) \rightarrow \alpha_0 \Phi(\mathbf{r}'_1, \mathbf{r}'_2) \Phi^*(\mathbf{r}_1, \mathbf{r}_2), \quad (13)$$

where  $\alpha_0$  is the largest eigenvalue of  $O(N)$ , and  $\Phi$  the corresponding eigenfunction<sup>2</sup>.

Applying the factorization property of the ODLRO hypothesis to both sides of the relation (11), we obtain, in the ODLR limit,

$$\begin{aligned} & \Phi(\mathbf{r}'_1, \mathbf{r}'_2) \Phi^*(\mathbf{r}_1, \mathbf{r}_2) = \\ & \exp\left\{\frac{ie}{c\hbar}[\chi_{\mathbf{a}}(\mathbf{r}'_1) + \chi_{\mathbf{a}}(\mathbf{r}'_2) - \chi_{\mathbf{a}}(\mathbf{r}_1) - \chi_{\mathbf{a}}(\mathbf{r}_2)]\right\} \Phi(\mathbf{r}'_1 - \mathbf{a}, \mathbf{r}'_2 - \mathbf{a}) \Phi^*(\mathbf{r}_1 - \mathbf{a}, \mathbf{r}_2 - \mathbf{a}), \end{aligned} \quad (14)$$

which implies<sup>5</sup>

$$\Phi(\mathbf{r}_1, \mathbf{r}_2) = f(\mathbf{a}) \exp\left\{\frac{ie}{c\hbar}[\chi_{\mathbf{a}}(\mathbf{r}_1) + \chi_{\mathbf{a}}(\mathbf{r}_2)]\right\} \Phi(\mathbf{r}_1 - \mathbf{a}, \mathbf{r}_2 - \mathbf{a}), \quad (15)$$

where  $f(\mathbf{a})$  is a possible displacement dependent, but position independent, phase factor<sup>6</sup>, with  $f(0) = 1$ . Thus, the function  $\Phi$ , which is to be identified with the Ginsburg-Landau wave function, at different space points is related by a phase factor. Repeating successively for infinitesimal displacements along a closed path  $C$ , and picking up products of phase factors and the path-dependent f-factors, we obtain from (15) the relation

$$\Phi(\mathbf{r}_1, \mathbf{r}_2) = F(C) \exp\left\{i\frac{2e}{c\hbar} \oint_C \mathbf{A}(\mathbf{r}) \cdot d\mathbf{r}\right\} \Phi(\mathbf{r}_1, \mathbf{r}_2), \quad (16)$$

where  $F(C)$  represents the product of the f-factors, and use has been made of the single-valuedness of the function  $\Phi(\mathbf{r}_1, \mathbf{r}_2)$ , a consequence of the single-valuedness of the density matrix.

### Meissner Effect

We now use (15) to prove the Meissner effect. In a simply connected region we apply (15) to two sequences of successive displacements<sup>7</sup>, first  $\mathbf{a}$  followed by  $\mathbf{b}$ , and then  $\mathbf{b}$  followed by  $\mathbf{a}$ . Comparing the two resulting relations, we obtain

$$\begin{aligned} & \exp\left\{\frac{ie}{c\hbar}[\chi_{\mathbf{a}}(\mathbf{r}_1) + \chi_{\mathbf{b}}(\mathbf{r}_1 - \mathbf{a}) + \chi_{\mathbf{a}}(\mathbf{r}_2) + \chi_{\mathbf{b}}(\mathbf{r}_2 - \mathbf{a})]\right\} = \\ & \exp\left\{\frac{ie}{c\hbar}[\chi_{\mathbf{b}}(\mathbf{r}_1) + \chi_{\mathbf{a}}(\mathbf{r}_1 - \mathbf{b}) + \chi_{\mathbf{b}}(\mathbf{r}_2) + \chi_{\mathbf{a}}(\mathbf{r}_2 - \mathbf{a})]\right\}. \end{aligned} \quad (17)$$

Namely,

$$\exp\left\{i\frac{2e}{c\hbar} \oint_{(\mathbf{a} \times \mathbf{b})} \mathbf{A}(\mathbf{r}) \cdot d\mathbf{r}\right\} = 1, \quad (18)$$

where the loop path of integration is a parallelogram formed by  $\mathbf{a}$  and  $\mathbf{b}$ . Therefore,

$$\mathbf{B} \cdot (\mathbf{a} \times \mathbf{b}) = n \frac{ch}{2e}, \quad n = \text{integers}. \quad (19)$$

While  $\mathbf{a}$  and  $\mathbf{b}$  can be continuously changed, the right-hand side of (19), being proportional to an integer, can not. Consistency then requires

$$\mathbf{B} = 0. \quad (20)$$

This is the property of the Meissner effect.

### Flux Quantization

Next, we use (15) and (16) to prove the quantization of flux, when the loop  $C$  is not simply-connected, like a path winding around a tunnel region inaccessible to the electrons.

As we have already proved the Meissner effect, namely  $\mathbf{B} = 0$ , it is clear from (16) that  $F(C) = 1$  if  $C$  is simply-connected. To prove the quantization of flux, we need to prove  $F(C) = 1$  for multiply-connected loops  $C$ .

Now, the factor  $f(\mathbf{a})$  in (15) is independent of  $\mathbf{r}$ . This implies that the property of  $F(C)$  is independent of the location of the loop  $C$ . Corresponding to a multiply-connected loop  $C$ , we can consider a loop of the same shape in a simply-connected region, for which  $F(C) = 1$ . It follows that  $F(C) = 1$  also for  $C$  in a multiply-connected region. In general, therefore,

$$F(C) = 1 \quad (21)$$

for any loop path  $C$ . (16) then becomes

$$\Phi(\mathbf{r}_1, \mathbf{r}_2) = \exp\left\{i \frac{2e}{c\hbar} \oint_C \mathbf{A}(\mathbf{r}) \cdot d\mathbf{r}\right\} \Phi(\mathbf{r}_1, \mathbf{r}_2). \quad (22)$$

Consequently, there is the quantization condition

$$\oint_C \mathbf{A}(\mathbf{r}) \cdot d\mathbf{r} = n \frac{ch}{2e}, \quad n = \text{integers}, \quad (23)$$

The loop path  $C$  may or may not be simply-connected. When  $C$  in (23) is simply-connected, it can be altered infinitesimally and continuously, implying, again, the Meissner

effect. When  $C$  is not simply-connected, like a path winding around an inaccessible tunnel, the path can not be deformed across this inaccessible region. Then the quantization condition (23) implies that the total flux<sup>8</sup> passing through the loop  $C$  can only have quantized values. We note that in our idealized considerations the loop path  $C$  is away from the boundaries.

We can prove  $F(C) = 1$  more explicitly by observing some simple properties of  $f(\mathbf{a})$ :

1. As  $f(\mathbf{a})$  is independent of  $\mathbf{r}$ , we can choose to explore the properties of  $f(\mathbf{a})$  by considering displacements in regions where  $\mathbf{B} = 0$ .

2. In any region of  $\mathbf{B} = 0$ , consider the succession of two infinitesimal displacements  $\mathbf{a}$  and  $\mathbf{b}$ , as well as the displacement  $\mathbf{a} + \mathbf{b}$ . Because  $\mathbf{B} = 0$  in the region enclosed by the three vectors  $\mathbf{a}$ ,  $\mathbf{b}$  and  $\mathbf{a} + \mathbf{b}$ , it is clear from (15) that

$$f(\mathbf{a} + \mathbf{b}) = f(\mathbf{a})f(\mathbf{b}). \quad (24)$$

(24) offers another more explicit proof of (21). Recall that  $F(C)$  is the product of the  $f$ -factors along the loop path. Hence, for a closed loop, on account of (24), we again obtain

$$\begin{aligned} F(C) &= f(\mathbf{a})f(\mathbf{b})f(\mathbf{c}) \cdots \\ &= f(\mathbf{a} + \mathbf{b} + \mathbf{c} + \cdots) \\ &= f\left(\oint_C d\mathbf{r}\right) = f(0) = 1. \end{aligned}$$

### Inclusion of Lattice Potential

The lattice potential  $V_c(\mathbf{r})$ , which is not invariant under arbitrary displacements, has so far been omitted in our consideration. We now consider the case when it is included.

Let the symmetry of the lattice be defined by three basic lattice vectors:  $\mathbf{e}_1$ ,  $\mathbf{e}_2$ ,  $\mathbf{e}_3$ . The lattice potential  $V_c(\mathbf{r})$  is then invariant under the space displacement

$$\mathbf{r}_j \rightarrow \mathbf{r}_j - \Delta\mathbf{r}, \quad (25)$$



if<sup>9</sup>

$$\Delta \mathbf{r} = n_1 \mathbf{e}_1 + n_2 \mathbf{e}_2 + n_3 \mathbf{e}_3, \quad n_1, n_2, n_3 = \text{integers}. \quad (26)$$

Under such displacements, the total Hamiltonian is invariant, except for an induced gauge term in the vector potential, like that in (4). This is exactly like the situation we had when the lattice potential was omitted, except, of course, that the displacements have to be restricted to those in (26). All previous arguments apply, leading to the conclusion, similar to (19), that

$$\mathbf{B} \cdot (\mathbf{e}_j \times \mathbf{e}_k) = n_{jk} \frac{ch}{2e}, \quad n_{jk} = \text{integers} \quad (27)$$

which, in turn, implies that

$$\mathbf{B} = \frac{1}{V} \frac{ch}{2e} \sum_j n_j \mathbf{e}_j, \quad n_j = \text{integers} \quad (28)$$

where

$$V = \mathbf{e}_1 \cdot (\mathbf{e}_2 \times \mathbf{e}_3).$$

Namely, the magnetic field  $\mathbf{B}$ , if non-vanishing, inside the material can only be along a discrete set of directions. Now, following Sewell<sup>3</sup>, we make the assumption that  $\mathbf{B}$  is along the direction of the external applied field. As the direction of the external field can be arbitrarily chosen, the magnetic field  $\mathbf{B}$  inside the material has to vanish, if not to contradict (28), giving rise to the Meissner effect.

In the above proof we have made the assumption that the direction of  $\mathbf{B}$  inside the material lies along the direction of the external field. We can, however, present another argument without having to make this assumption. Again, consider the requirement imposed by (28). Typical lattice dimension of superconducting materials is on the order of  $1\text{\AA}$ . The smallest value of  $B$ , if not zero, has to be on the order of

$$B \sim (1\text{\AA})^{-2} \frac{ch}{2e} \sim 10^9 G, \quad (29)$$

according to (28). This is many orders stronger than the typical critical field strengths of less than  $10^3 G$ , beyond which the superconducting state no longer exists, for superconductors. Therefore, to be consistent with (28), the only possible value of  $\mathbf{B}$  is zero for a superconducting state.

The same reasoning that lead to the quantization condition (23) applies, and leads to

$$\sum_C \mathbf{A}(\mathbf{r}) \cdot \Delta \mathbf{r} = n \frac{ch}{2e}, \quad n = \text{integers}, \quad (30)$$

where loop path  $C$  consists of a succession of displacements of the type specified by (26). This quantization condition, when applied to multiply-connected paths, implies the phenomenon of flux quantization.

## Discussion

We conclude with a few remarks.

1. The function  $\Phi(\mathbf{r}, \mathbf{r})$  corresponds to the Ginsburg-Landau wave function<sup>10</sup>. The successive displacements taken in arriving at equation (16) clearly demonstrates the origin of the characteristics of the superconducting state: Coherence of phases is responsible for the properties of flux quantization and the Meissner effect. It is also made clear by the proof given in this note that these twin properties of superconductivity have a common origin, an understanding perhaps not sufficiently emphasized in usual presentations.

2. The aspect in the hypothesis of ODLRO that has been made use of in the proof of flux quantization and the Meissner effect is the factorization property<sup>2</sup> of  $\rho_2$  as in (13), which is responsible for the coherence of phases in  $\Phi(\mathbf{r}_1, \mathbf{r}_2)$ . The ODLRO hypothesis implicitly assumes that in the sum in (12) there is only one term, having a large eigenvalue on the order of  $O(N)$ , that survives in the ODLR limit.

## Acknowledgement

One of the authors (HTN) has benefited from discussions with A. S. Goldhaber, J. K. Jain, V. Korepin, T. K. Ng and W. Weisberger. We would also like to thank Prof. C. N. Yang, from whom they have learned over the years the fundamentals of superconductivity. Gang Su is grateful to Prof. C. N. Yang and ITP for the kind hospitality during his visit, and the Charles C. K. Yeung fellowship through the CEEC of SUNY at Stony Brook for support. Two of the authors (GS and BHZ) are supported in part by the National Natural Science Foundation of China.

## Footnotes

<sup>1</sup>O. Penrose, Phil. Mag. **42**, 1373(1951); O. Penrose and L. Onsager, Phys. Rev. **104**, 576(1956).

<sup>2</sup>C. N. Yang, Rev. Mod. Phys. **34**, 694(1962).

<sup>3</sup>G. L. Sewell, J. Stat. Phys. **61**, 415(1990).

<sup>4</sup> In proving the flux quantization, we shall need to consider the case where there are regions inaccessible to the electrons, like a tunnel. In these inaccessible regions, the magnetic field does not necessarily vanish. In order to guarantee the single-valuedness of phase  $\chi_{\mathbf{a}}(\mathbf{r}_j)$  in (7), we need to restrict to infinitesimal  $\mathbf{a}$ , so that  $\phi(\mathbf{r}_j - \mathbf{a}) - \phi(\mathbf{r}_j)$  is single-valued, although  $\phi(\mathbf{r}_j - \mathbf{a})$  and  $\phi(\mathbf{r}_j)$  individually are not.

<sup>5</sup>Here  $\Phi(\mathbf{r}_1, \mathbf{r}_2)$  has been renormalized, i.e., a factor  $\sqrt{\alpha_0}$  has been absorbed into it.

<sup>6</sup>We thank W. Weisberger for pointing out the possible existence of this displacement-dependent, but  $\mathbf{r}$ -independent phase factor.

<sup>7</sup>The consideration parallels Sewell's, in Ref. 3, in his derivation of the Meissner effect.

<sup>8</sup>The total flux includes the flux through the tunnel as well as the flux of the penetrated magnetic field into the boundary layer surrounding the tunnel.

<sup>9</sup>For reasons explained in footnote<sup>4</sup>, we only consider  $\Delta\mathbf{r}$  which is infinitesimal on macroscopic scale. Thus, the integers  $n_1$ ,  $n_2$  and  $n_3$  can not be large.

<sup>10</sup>It is the factorization property that was assumed by L.P. Gor'kov [JETP **7**, 505(1958)] to derive his effective equations for superconductivity.